

To Cite:

Sharma S. Utilization of non-timber forest products in Panchamul-1, Andhikhola Rural Municipality, Syangja district, Nepal. *Discovery*, 2021, 57(307), 578-596

Author Affiliation:

Kathmandu Forestry College, Institute of Forestry-
Tribhuvan University, Kathmandu, Nepal

Correspondence to:

Email: surajsharma83@yahoo.com

Peer-Review History

Received: 23 May 2021

Reviewed & Revised: 24/May/2021 to 27/June/2021

Accepted: 28 June 2021

Published: July 2021

Peer-Review Model

External peer-review was done through double-blind method.



© The Author(s) 2021. Open Access. This article is licensed under a Creative Commons Attribution License 4.0 (CC BY 4.0), which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

Utilization of non-timber forest products in Panchamul-1, Andhikhola Rural Municipality, Syangja district, Nepal

Suraj Sharma✉

ABSTRACT

The present paper documents the utilization of Non-Timber Forest Products in Panchamul-1, Andhikhola Rural Municipality, Syangja district of Nepal. It also determines the informant's knowledge and perception of medicinal plants, fodder plants, and wild edible plants utilized in the study area. The information regarding ethnobotany, fodder plants, and wild edible plants was collected through a pre-structured questionnaire survey, Key Informant Interview (KII), and Focus Group Discussion (FGD). Data based on ethnobotany were analyzed through Informant Consensus Factor (ICF), Fidelity Level (FL %), and use value (UV). Quality of fodder and the parts used of different Wild Edible Plants (WEPs) were based on informant knowledge and perception. In this study, a total of 111 plant species under 90 genera belonging to 59 families were recorded in Panchamul. Of which 62 species were used treating 64 ailments, where Cough/Cold ailment category has the highest ICF value (0.67) with 38 use reports and 13 plant species. Whereas, the least agreement was seen in the ailment category of Cuts/Wounds which have the lowest ICF value (0.043) with 15 use reports, 9 plant species. *Asparagus racemosus* has the highest FL (100%) used for Gastrointestinal complaints followed by *Halenia elliptica* (91.67%) used for fever, and *Lindera neesiana* (45.61%) has the lowest FL to cure Skeleto-muscular system problems. 12 fodder plants were identified as best fodder among 31 plants *viz.* *Artocarpus lacucha*, *Arundinaria falcate*, *Bauhinia purpurea*, *Dendrocalamus hamiltonii*, *Erythrina stricta*, *Ficus lacor*, *Ficus nemoralis*, *Ficus semicordata*, *Ficus subincisa*, *Premna bengalensis*, *Premna latifolia*, and *Quercus semecarpifolia*. Wild edible plants used in Panchamul-1 were mainly fruits (16) followed by leaves (11), tuber (06), etc. and these wild edible plants were mostly used for raw fruits (16), followed by vegetables (13), pickles (8) boil (5), spices (3), and jam (2). Based on ICF values, it is seen that there was a very high agreement in the usage of medicinal plants, which belongs to the cough/cold ailment category. Use value and fidelity level also indicate the most desired medicinal plant species used by the local inhabitants. Also, the informant's knowledge and perception regarding medicinal plants, fodder plants, and wild edible plants were well documented and it needs more in-depth research and conservation priority to those preferred plant, though these species were not seen harvested for trade in the study area.

Keywords: Edible plants, fodder plants, NTFPs, Medicinal Plants.

1. INTRODUCTION

Nepal is very rich in cultural as well as biological diversity, though a small country with an area of 147516 Km² hosts 10669 species of flora (Hara and William, 1979; Hara et al., 1978; Chaudhury, 1998). Altogether 336 species of flora are endemic to Nepal (Joshi and Joshi 1991; Shrestha and Joshi 1996; Chaudhury, 1998). Also, it stands in the 10th position in having the highest diversity of flowering plants i.e. 6501 numbers in total (DPR, 2001). Nepal with such a rich floral diversity is not yet able to harness the benefits from the existing resource base (Subedi, 1997). In Nepal, both in-situ and ex-situ conservation and management of plant biodiversity are being practiced (Chaudhary 1998; Chaudhary et al. 2016) with very strong forest governance policies (MoEF, 2018). In Nepal 700 medicinal plants, 440 wild foods, 30 spices, and the other 71 are fibers yielding (Subedi et al., 2014).

NTFP is defined as “all products derived from biological resources found on forest land, but not including timber or fuelwood” (Wong, 2002). Depending on the objectives NTFPs are defined in either way as: ‘non-timber forest product’, ‘non-wood forest product’, ‘wild products’, ‘natural products’, ‘non-timber forest and grassland products’, ‘minor forest products’, ‘secondary forest products’, ‘byproducts of the forest’, ‘non-traditional forest resources’, ‘non-timber trees product’ or ‘agroforestry products’ (FAO, 1999). Rural communities of the developing country use NTFPs in a huge amount either for medicine, fodder, fertilizers, foods like fruits, nuts, honey, etc. construction materials, cosmetics, natural dyes, tannin, gums, aromatic oils, spices, edible oils, horns, decorative articles, bones, etc. These products can be derived from various sources such as plants (leaves, root, flower, stem, etc.), animals, and other non-living components of the ecosystem (Akanni, 2013). Sustainable utilization of NTFPs and its promotion could lead to poverty reduction, biodiversity conservation (FAO, 1995; Shiva and Verma, 2002; Golam et al., 2008), and hence maintaining the forest environmental services and biological diversity (Ros-Tonen, 2000). While some (Peters et al., 1989) suggest that NTFP extraction is financially and ecologically sustainable, others point to its adverse social and ecological consequences (Arnold and Perez, 2001; SCBD, 2001). In developing countries like Nepal, any conservation program should be based on a broad understanding of landscape ecology (dynamics) and its drivers of social, economic, and ecological changes, which will not only replace but complement the conventional biodiversity conservation practice. Thus, the socio-economic well-being of the indigenous groups (Sayer 2009, Kremen and Merenlender, 2018).

Although, NTFPs, harvested for subsistence purposes (local use), exhibits little pressure. But commercially demanded NTFPs species are over-harvested (unsustainable harvesting) and it could lead to extinction threat. Often, NTFP species are overexploited because of their higher market values (Edward, 1994; Karki, 1996; Sharma, 1996), an unclear definition of property rights, a lack of knowledge on conservation, and increasing market demand (Subedi and Bhattarai, 1998). The Government of Nepal has endorsed acts, policies, plans, and legislative measures to regulate the collection, production, trade, processing, and marketing of NTFPs. However, the rules aren't regulated when NTFPs are harvested and traded. This study will emphasize the use-value and its importance of NTFPs the medicinal plants (ailment categories), fodder plants, and wild edible plants in the communities and stimulate the interest of the local inhabitants in the sustainable use and management of NTFPs.

2. MATERIALS AND METHODS

Study Area

Andhikhola is a Rural Municipality in Syangja Districts in the Gandaki Province of central Nepal. The total population of Aandhikhola is 16,589 with a total area of 69.61 km² (NPHC, 2011). It lies between 28° 6' N Latitude and 83° 45' E Longitude. Panchamul (Fig. 1) Ward-1 of this rural municipality comprises a total of 1079 households with a total of 6458 individuals (3175 female and 3283 male) as per the Panchamul ward office electoral roll report. The total area of Panchamul -1 is approximately 12.71 km² with (708 Acre) 2.87 km² forest cover.

Data Collection

The survey techniques and inventory techniques (Martin, 1995; Rastogi et al., 1998; Cunningham, 2001; Ghimire et al., 1999, 2000, 2001) were used for field data collection. Primary data were collected with an amalgam of survey methods including reconnaissance survey, participatory techniques- Focus Group Discussions (FGD), Key Informant Interview (KII), and formal & informal interviews. A simple random and purposive sampling method is used to administer the semi-structured questionnaire survey of households and on-site observations. The inventory technique by Transect walk survey was used for species identification by their local names and parts used, based on key informant's knowledge (Ghimire, et al., 1999, 2000, 2001). The NTFPs were also identified taking help of experts, taxonomist, and standard literature (Hara et al., 1978, 1982; Hara and Williams, 1979; Polunin and Stainton, 1984; Stainton, 1988; Press et al., 2000; Lama et al., 2001; DPR, 2001; Zheng-yi et al., 1994).




Figure: 1. Study Area

A formula based on (Cochran, 1977) was used to estimate the sample size (n) to administer the pre-structured questionnaire at a 95% confidence level.

$$\text{Sample Size } (n) = \frac{N \cdot z^2 \cdot P(1-P)}{N \cdot d^2 + z^2 \cdot P(1-P)}$$

Where ' n ' is sample size with finite population correction, N is the population size, ' z ' is z statistic for a level of confidence, ' P ' is the expected proportion, and ' d ' is the precision.

Data Analysis

The qualitative data were analyzed and described in the text. The quantitative data were analyzed in SPSS 16 and MS excel 2013. The results were presented in a table, pi-charts, and bar diagrams.

Medicinal Used Value (UV)

Plants species used as medicine and their relative importance was estimated with the help of formula on use value (UV) for i^{th} species.

$$UV_i = \frac{\sum U_i}{N_i}, \text{ (Phillips et al., 1994)}$$

Where U_i is the number of use-reports cited by each informant for a given plant species i and N_i is the total number of informants interviewed for a given plant species i . Use values are high when there are many use-reports for a plant and low when there are few reports related to its use.

Informant Consensus Factor (ICF)

Informant Consensus Factor (ICF) was used to estimate the general agreement (i.e. the relative homogeneity of use-reports) in the usage of medicinal plants in various ailments categories in the study area. The formula used to calculate ICF for the j^{th} species is:

$$ICF_j = \frac{Nur_j - Nt_j}{Nur_j - 1}, \text{ (Trotter and Logan, 1986; Heinrich et al., 1998)}$$

Where Nur_j is the number of use-reports in each ailment category j and Nt_j is the total number of taxa used in each ailment category j by all informants. A higher value indicates a high rate of agreement, a low value indicates a low degree of agreement (Gazzaneo et al., 2005).

Fidelity level (FL)

In the study area, fidelity level (FL) was calculated to determine the frequently used plant species (i^{th}) by the local people for treating a specific ailment category (j) using the following formula:

$$FL_{ij} (\%) = \frac{N_{pij}}{N_i} \times 100, \text{ (Friedman et al., 1986)}$$

Where N_{pij} is the number of use-reports cited for a given plant species i for a particular ailment category j and N_i is the total number of use-reports cited for any given species i . The highest FL_{ij} value of plant species is considered a highly preferred species for ailment category j . Spearman's correlation coefficient (data were not normally distributed) was used to determine the correlations between use-value (UV) and fidelity level (%) for the medicinal plants documented in this study.

3. RESULTS AND DISCUSSIONS

A pre-structured questionnaire survey among 148 individuals and 28 key informants (18 Male and 10 Female) between ages 25 to 75 years was administered (Table 1). About 35% (10) were illiterate, 25% (7) had primary education, 32% (9) had secondary education, whereas, only 8% (2) were graduates. Among the 28 individuals selected key informants comprised of local healers (Brahmins) and farmers (Dalits, Gurungs, and Brahmins).

Table 1: Gender and Age Class of the Key Informants						
Gender	Informants Age class in Years					
	≤30	31-40	41-50	51-60	61-70	≤75
Female	0	1	3	3	2	1
Male	1	3	4	3	4	3
Total	1	4	7	6	6	4

Table 2: Medicinal Plants used by the resident of Panchamul, Andhikhola, Syangja					
Botanical Name (Family)	Vernacular Name	Medicinal Plants		UV _i	Similar Findings (references)
		Parts	Treatment		
<i>Achyranthes bidentata</i> Bl. (Amaranthaceae)	Datiwin (Herb)	Ro, St	Toothache, Evil eyes	0.75	Thapa (2012)
<i>Aconitum spicatum</i> Staf. (Ranunculaceae)	Bikh (Herb)	Ro, L	Analgesic	0.43	Lama et al. (2001), Manandhar (2002) Joshi and Joshi (2001), and Baral and Kurmi (2006)
<i>Acorus calamus</i> L. (Araceae)	Bojho (Herb)	Ro	Toothache, Headache, Cough, Fever, Anti-fungal	0.64	Joshi and Joshi (2001)
<i>Amaranthus spinosus</i> L. (Amaranthaceae)	Lunde (Herb)	Ro, L	Over body heat	0.43	Thapa (2012)
<i>Anemone polyanthes</i> D. Don (Ranunculaceae)	Avijalo (Herb)	WP	Indigestion, Cough, Cold	0.89	Ghimire et al. (2001)
<i>Artemisia vulgaris</i> L. (Asteraceae)	Titepati (Shrub)	L	Antiseptic, Antibacterial and Antimicrobial	0.75	Rysdyk (2019)
<i>Asparagus racemosus</i> Wild. (Asparagaceae)	Kurilo (Climber)	Ro	Stomachache, constipation, anxiety, Breast milk production	0.60	Pizzorno, et al. (2015), Hechtman (2018), Goyal et al. (2003)

<i>Berberis aristata</i> DC. (Berberidaceae)	Chutro (Shrub)	Fr, Br, L	Eye problems, Fever, Jaundice	0.25	Ghimire et al. (2001), Rajbhandari (2001), Lama et al. (2001), Manandhar (2002), Baral and Kurmi (2006) and Kunwar et al. (2006)
<i>Bergenia ligulata</i> (Wall.) (Saxifragaceae)	Phakanbed (Herb)	Ro, Rh	Fever, Anti-diabetic, Lowers Blood Pressure, Kidney stone.	0.18	Gurav and Gurav (2014)
<i>Calotropis gigantea</i> (L.) (Apocynaceae)	Ankh (Shrub)	Lx	Blood Clot, Fracture	0.18	Bhattarai, et al., (2009)
<i>Choerospondias axillaris</i> (Roxb.) B. L. Burt & A. W. Hill (Anacardiaceae)	Lapsi (Tree)	Fr, Br	Anti-oxidant, Secondary burn	0.36	Wang et al. (2008), Nguyen et al. (1996).
<i>Centella asiatica</i> (L.) Urban (Apiaceae)	Ghodtapre (Herb)	WP	Anxiety, Common cold and flu, Diarrhea, Fatigue, Hepatitis, Indigestion, Jaundice, Tonsillitis, Urinary tract infection, wound healing	1.00	Shrestha and Dhillion, (2003), Tamang, (2003), Bhattarai, et al., (2009), Joshi, et al., (2011), Thapa, (2012)
<i>Cheilanthes dalhousiae</i> Hook. (Pteridaceae)	Dankasnu (Fern)	L	Gastritic	0.31	Thapa (2012)
<i>Cinnamomum tamala</i> (Buch.- Hum.) Ness & Eberm. (Lauraceae)	Tejpatta (Tree)	L	Stomach Gas, Stomachache, Diarrhea, Allergy, Cough and Cold	0.57	Hasan et al. (2013)
<i>Crateva unilocularis</i> Buch.- Ham. (Capparaceae)	Siplikaan (Tree)	L	Stomach ache, Bad cholesterol, Blood pressure, Fever, Urinary Problems	0.39	Manandhar (2002)
<i>Curcuma caesia</i> Roxb. (Zingiberaceae)	Kalo Haledo (Herb)	Rh	Dysentery, Back pain	0.50	Thapa (2012)
<i>Curcuma zedoaria</i> (Christm.) <i>Roscoe</i> (Zingiberaceae)	Seto Haledo (Herb)	Rh	Dysentery, Back Pain	0.57	Thapa (2012)
<i>Drynaria propinqua</i> (Wall. ex Mett.) J.Sm. ex Bedd. (Polypodiaceae)	Kammari Laharo (Climber)	St, L	Bone Fracture	0.75	Chang et al. (2007)
<i>Eupatorium adenophorum</i> Spreng. (Compositae)	Banmara (Shrub)	L	Antiseptic, Blood Coagulant	0.93	Thapa (2012)
<i>Euphorbia hirta</i> L. (Euphorbiaceae)	Dodhe Jhar (Herb)	Ro	Antiseptic, Blood Coagulant, Skin Burn, Piles	0.78	Thapa (2012)
<i>Euphorbia royleana</i> Boiss. (Euphorbiaceae)	Siudi (Shrub)	Lx	Skin diseases	0.28	
<i>Ficus semicordata</i> Buch.-Ham. ex Sm. (Moraceae)	Khanyu (Tree)	L	Scabies	0.43	Thapa (2012)
<i>Fragaria nubicola</i> Lindl. ex Lacaita (Rosaceae)	Bhui Aiselu (Herb)	Fr, L	Skin Rashes (Herpes zoste)	0.57	Thapa (2012)
<i>Girardinia diversifolia</i> a (Link) Friis (Urticaceae)	Chalne Sisnu (Herb)	Ro	Swollen body, Internal injury, Blood purification	0.25	Manandhar (2002), Kunwar et al. (2006) and Ghimire et al. (2001)
<i>Gonostegia hirta</i> (Bl.) Miquel	Chiple Ghas	WP	Dislocated Bone	0.28	Thapa (2012)

(Urticaceae)	(Herb)				
<i>Halenia elliptica</i> D. Don (Gentianaceae)	Tite (Herb)	L, St	Fever, Headache, Cough And Cold	0.78	Rajbhandari (2001), Pohle (1990), Chophel (1993), Dawa (1993)
<i>Houttuynia cordata</i> Thunb. (Saururaceae)	Gandhe (Herb)	WP	Stomachache, Gastric, Boost appetite	0.68	Yang and Jiang (2009)
<i>Jatropha curcus</i> L. (Euphorbiaceae)	Sajiwan (Shrub)	Lx	Toothache, Anti-microbial	0.25	Patil et al., (2013)
<i>Juniperus indica</i> Bertol (Cupressaceae)	Dhupi (Shrub)	L, Fr	Cough and Cold, Skin Diseases, Incense	0.28	Phole (1990), Joshi and Joshi (2001), Baral and Kurmi (2006) and Kunwar et al. (2006)
<i>Lagerstroemia parviflora</i> Roxb. (Lythraceae)	Dhaiyanro (Tree)	F	Dysentery (Blood stool)	0.18	
<i>Lindera neesiana</i> (Wall. ex Nees) Kurz (Lauraceae)	Siltimur (Tree)	S	Cough, Cold, Headache	0.93	Thapa (2012)
<i>Lycopodium clavatum</i> L. (Lycopodiaceae)	Nagbeli (Herb)	WP, Sp	Cough, Rheumatism, Gastric, Constipation, Fever	0.14	Bharadwaj and Mishra (2018)
<i>Lyonia ovalifolia</i> (Wall.)Drude. (Ericaceae)	Angeri (Shrub)	L, B	Skin disease	0.21	Manandhar (2002)
<i>Mirabilis jalapa</i> L. (Nyctaginaceae)	Seto Malati (Herb)	Ro	Menstrual Problem, Dysentery, Headache, Earache	0.38	Boullard (2001)
<i>Morus alba</i> L. (Moraceae)	Kafal (Tree)	Ro	Intestinal worm	0.11	Thapa (2012)
<i>Myrica esculenta</i> Buch.-Ham. ex. D. Don (Myricaceae)	Ban Kafal (Tree)	Fr, Br, L	Asthma, Cough, Chronic Bronchitis, Ulcers, Constipation, Anemia, Fever, Diarrhea, And Ear, Nose, And Throat Disorders.	0.32	Kumar and Rana (2013), Sood and Shri (2018) Shrestha and Dhillion, (2003), Tamang and Sedai (2016)
<i>Nardostachys grandiflora</i> DC. (Valerianaceae)	Jatamansi (Herb)	Rh	Headache, Cough, Cold, Fever,	0.71	Ghimire et al. (2001), Kunwar et al. (2006)
<i>Opuntia monacantha</i> Haw. (Cactaceae)	Paate Siundi	WP	Burns, Wounds, Indigestion, Diabetes	0.78	Hasan (2018)
<i>Oxalis corniculata</i> L (Oxalidaceae)	Amilo Jhar (Herb)	WP	Sinus, Stomach Problems, Appetite	0.93	Thapa (2012)
<i>Parmelia nepalensis</i> Tayl. (Parmeliaceae)	Jhyau (Lichen)	WP	Menstrual Disorder	0.43	NEHHPA
<i>Periploca calophylla</i> (Wight) Falc. (Asclepiadaceae)	Sikari Lahara (Climber)	WP	Fractured Bone, Body ache, Backache	0.28	Adhikari, et al. (2019)
<i>Phyllanthus emblica</i> L. (Euphorbiaceae)	Amla (Tree)	Fr	Cough, Sore Throat, Constipation, Diarrhea, Gastritis	0.96	Bhattarai, et al., (2009), Luitel, et al. (2014), Tamang and Sedai (2016)
<i>Pogostemon plectranthoides</i> Desf. (Lamiaceae)	Rudilo (Herb)	Ro,	Fever	0.43	Dangol (2008)
<i>Polypodium vulgare</i> L. (Polypodiaceae)	Bikhfage (Fern)	Rh	Constipation, Diarrhea	0.28	Camps et al. (1990)
<i>Prunus cerasoides</i> D. Don (Rosaceae)	Paiyun (Tree)	Br	Barks are cut into pieces, boiled and decoction is	0.25	Manandhar (2002)

			applied externally on swollen and burnt body parts.		
<i>Psidium guajava</i> L. (Myrtaceae)	Amba (Tree)	L, Br	Anthelmintic, Stomachache, Blood stool	0.78	Bhattarai, et al., (2009), Malla and Chhetri (2009), Luitel, et al. (2014), Tamang and Sedai (2016)
<i>Rhododendron arboretum</i> Sm. (Ericaceae)	Lali Guras (Tree)	L, Br	Headache, Jaundice, Diabetes, liver disorders, and Intestinal worms	0.64	Rawat et al. (2017)
<i>Rosa serisea</i> Lindl. (Rosaceae)	Bhote Gulab (Herb)	Fl	Headache	0.21	Manandhar (2002)
<i>Rubia manjith</i> Roxb. ex Fleming (Rubiaceae)	Majhitho (Climber)	Ro	Fever, Chest Pain, Dysentery, Rheumatism	0.14	Ghimire et al. (2001), Manandhar (2002)
<i>Rubus ellipticus</i> Sm. (Rosaceae)	Aniselu (Shrub)	WP	Gastric, Sinusitis	0.43	Shrestha and Dhillion, (2003), Thapa (2012), Luitel, et al. (2014), Tamang and Sedai (2016), Joshi, et al., (2011)
<i>Rumex nepalensis</i> Spreng. (Polygonaceae)	Halhale (Herb)	Ro, L	Constipation, Diarrhea, Vegetable/Pickle	0.39	Manandhar (2002), Kunwar et al., (2006)
<i>Selinum tenuifolium</i> Wall. ex C.B. Clarke (Apiaceae)	Bhutkase (Herb)	WP, Ro	Traditional Magico-Religious beliefs	0.14	Umberto, (2012)
<i>Solanum capsicoides</i> All. (Solanaceae)	Kantakari (Herb)	Fr	Toothache,	0.25	Thapa (2012)
<i>Swertia chirayita</i> (Roxb. ex Fleming) Karsten. (Gentianaceae)	Chiraito (Herb)	WP	Cough, Cold, Fever, Malaria, and Headache	0.32	Rajbhandari (2001), Manandhar (2002), Rokaya et al., (2010)
<i>Tectaria coadunata</i> (Wall. ex Hook. & Grev.) C.Chr (Tectariaceae)	Kalo Nuro (Fern)	Rh	Dysentery	0.38	Thapa (2012)
<i>Tinospora sinensis</i> (Lour.) Merr. (Menispermaceae)	Gurjho (Climber)	St	Mensuration, Diabetes, Arthritis, Immunostimulant Diarrhea, Dysentery	0.78	Tamang, (2003), Bhattarai, et al., (2009)
<i>Trichosanthes tricusoidata</i> Lour. (Cucurbitaceae)	Indrayani (Tree)	Fr	Asthma, Earache, Rheumatism	0.43	Dubey et al. (2012).
<i>Tropaeolum majus</i> L. (Tropaeolaceae)	Barahmase (Climber)	L, Fl	Sinusitis	0.28	Thapa (2012)
<i>Urtica dioica</i> L. (Urticaceae)	Sisnu (Herb)	L, St, Ro	Chest Problems, Gastritis, Cuts & Wounds, Fractured & Broken Bones	0.78	Rajbhandari (2001), Ghimire et al. (2001), Manandhar (2002), Kunwar et al. (2006), Rokaya (2010)
<i>Valeriana jatamansi</i> Jones (Caprifoliaceae)	Sugandhawal (Herb)	WP	Dysentery, Diarrhea, Headache, Cough, and Indigestion.	0.32	Ghimire et al. (2001) and Kunwar et al. (2006)
<i>Vitex negundo</i> L. (Lamiaceae)	Simali (Shrub)	L, S, Ro	Menstrual Problem, Analgesic, Fever	0.25	Wan Hassan (2010), Kumar et al. (2018)
<i>Zanthoxylum armatum</i> DC (Rutaceae)	Timur (Shrub)	B, C, Fr, S	Toothache, Gastric	0.88	Thapa (2012), Hasan et al. (2013)
C= Carpels, Br=Bark, B-Bulb, L=Leaves, WP=Whole Plant, St=Stem, Fr=Fruit, Ro=Root, Lx=Latex, S=Seed, Fl=Flower, Sp=Spores, UV _i = Use value (i th Species)					

In this study, a total of 111 plant species under 90 genera belonging to 59 families were recorded in Panchamul, treating 64 ailments. Out of the total plant species, 62 species were used as medicinal plants, 31 species were fodder plants, and 41 species were wild edible plant species. A total of 62 medicinal plants under 59 genera belonging to 42 families were recorded (Table 2). Among the recorded families Euphorbiaceae (4), Rosaceae (4), and Polygonaceae (3) had the highest species counts (Fig. 2). During the treatment process, it is seen that leaves (21) were the most used followed by roots (14), whole plants (11), fruits (9), bark and rhizomes (6), etc. (Fig. 3). The results based on ICF (Informant Consensus Factor) showed that the value ranges from 0.43-0.67, where the Cough/Cold ailment category has the highest value i.e. ICF-0.67, 38 use reports, 13 plants species (*Centella asiatica* FL=73.68%, *Phyllanthus emblica* FL=71.05%, and *Lindera neesiana* FL=68.42% holds the highest Fidelity level) followed by Respiratory System Disorder category i.e. ICF- 0.66, 13 use reports, 5 plant species (*Trichosanthes tricusoidata* FL=92.31%, and *Myrica esculenta* FL=69.23% holds the highest Fidelity level) and then Circulatory System Disorders category i.e. ICF- 0.63, 42 use reports, 16 plant species (*Eupatorium adenophorum* FL=61.90%, and *Psidium guajava* FL=52.38% holds the highest Fidelity level). The least agreement was seen in ailment category of Cuts/Wounds i.e. ICF-0.043, 15 use reports, 9 plant species (*Centella asiatica* FL= 88.23%, *Opuntia monacantha* FL=68.18 %, and *Urtica dioica* FL=68.18 hold the highest Fidelity level) followed by Dermatological Infections category i.e. ICF- 0.44, 19 use reports, 11 plant species (*Artemisia vulgaris* FL=90.50%, *Euphorbia hirta* FL=86.36%, *Opuntia monacantha* FL=86.36%, and *Fragaria nubicola* FL=84.21% holds the highest Fidelity level) (Table 3). A similar study in Rasuwa district, Central Nepal (Upriety et al., 2010) reported disorders related to kidney problems, toothache, and ophthalmic problems showed the highest ICF of 1.0, and the lowest values were found for gastro-intestinal ailments category with an ICF of 0.53 and a study in Humla district, Western Nepal (Rokaya et al., 2010) reported Gastro-intestinal ailments category showed the highest ICF of 0.40, and the lowest values were found for Ophthalmological uses (ICF-00) followed by Respiratory system disorders (ICF-0.06). Similar studies from India and Thailand have reported much higher ICF values as mentioned in our study (Shristi et al., 2009; Rajkumar and Shivanna, 2009). Traditional use of specific medicinal plants and their knowledge were seen transferring orally to the young generation without any methodical process (Bussmann and Sharon, 2006) from generation to generation. These findings were supported by many studies (Joshi and Edington, 1990; Jain and Saklani, 1991; Shrestha and Dhillion, 2003; Rajkumar and Shivanna, 2009; Upriety et al., 2010; Panging and Sharma, 2017).




Fig 2: Families descriptions of available medicinal plants in Panchamul




Fig 3: Plants parts for treatment of various ailments

Table 3: Ailments (Categories) and informant consensus factor (ICF) and highest fidelity level				
Ailments (Categories)	Nur _j	Nt _j	ICF	FL (%)
Circulatory system disorders	42	16	0.63	
<i>Eupatorium adenophorum</i>				61.90
<i>Psidium guajava</i>				52.38
Cough/cold	38	13	0.67	
<i>Centella asiatica</i>				73.68
<i>Phyllanthus emblica</i>				71.05
<i>Lindera neesiana</i>				68.42
Cuts/wounds	15	9	0.43	
<i>Centella asiatica</i>				88.23
<i>Opuntia monacantha</i>				68.18
<i>Urtica dioica</i>				68.18
Dermatological infections	19	11	0.44	
<i>Artemisia vulgaris</i>				90.50
<i>Euphorbia hirta</i>				86.36
<i>Opuntia monacantha</i>				86.36
<i>Fragaria nubicola</i>				84.21
Ear, nose and throat (ENT) problems	21	10	0.55	
<i>Oxalis corniculata</i>				87.50
<i>Centella asiatica</i>				80.95
Fever	24	11	0.56	
<i>Halenia elliptica</i>				91.67
<i>Nardostachys grandiflora</i>				83.33
Gastro-intestinal ailments	83	46	0.46	
<i>Asparagus racemosus</i>				100
Genito-urinary ailments	17	7	0.62	

<i>Crateva unilocularis</i>				64.70
<i>Centella asiatica</i>				60.71
Respiratory system disorders	13	5	0.66	
<i>Trichosanthes tricusoidata</i>				92.31
<i>Myrica esculenta</i>				69.23
Skeleto-muscular system problems	57	31	0.46	
<i>Lindera neesiana</i>				45.61
Others/ Unclassified	9	5	0.50	
<i>Choerospondias axillaris</i>				90.00
<i>Berberis aristata</i>				77.78
<i>Jatropha curcus</i>				77.77
Total	338	164*		
* A taxon may be reported in more than one ailment category, FL= Fidelity level, Nur =Number of use-reports, Nt =Number of taxa				

Table 4: Fodder Plants used by the resident of Panchamul, Andhikhola, Syangja

Botanical Name (Family)	Vernacular Name	Quality of Fodder
<i>Aesandra butyracea</i> (Roxb.) Baehn (Sapotaceae)	Chiuree	Fair
<i>Albizia procera</i> Roxb. Benth. (Leguminosae)	Siris	Fair
<i>Artocarpus lacucha</i> Buch.-Ham. (Moraceae)	Badahar	Best
<i>Arundinaria falcata</i> Nees (Gramineae)	Neevalo	Best
<i>Bauhinia purpurea</i> L. (Leguminosae)	Tanki	Best
<i>Bauhinia variegata</i> L. (Leguminosae)	Koiralo	Good
<i>Brassaiopsis hainla</i> Buch.-Ham. Seem. (Araliaceae)	Chuletro	Good
<i>Buddleja asiatica</i> Lour. (Loganiaceae)	Phurse	Good
<i>Buddleja paniculata</i> Wall. (Loganiaceae)	Narayan Patti	Good
<i>Dalbergia sissoo</i> Roxb. (Leguminosae)	Sisou	Fair
<i>Dendrocalamus hamiltonii</i> Nees and Arn. (Gramineae)	Bans	Best
<i>Duabanga grandiflora</i> (Roxb.exDC.)Walp. (Lythraceae)	Lampate	Fair
<i>Engelhardia spicata</i> Leschen. Ex Blume (Juglandaceae)	Mahuwa	Fair
<i>Erythrina stricta</i> Roxb. (Leguminosae)	Phaledo	Best
<i>Ficus hispida</i> L.f. (Moraceae)	Tote	Good
<i>Ficus lacor</i> Buch.- Ham. (Moraceae)	Kavro	Best
<i>Ficus nemoralis</i> Wall. ex Miq. (Moraceae)	Dudhilo	Best
<i>Ficus semicordata</i> Buch.-Ham. ex Smith (Moraceae)	Khaneyu	Best
<i>Ficus subincisa</i> Buch.- Ham. Ex Smith (Moraceae)	Bidilno	Best
<i>Fraxinus floribunda</i> Wall. (Oleaceae)	Lankuri	Fair
<i>Melia azedarach</i> L. (Meliaceae)	Bakaino	Fair
<i>Morus australis</i> Poir. (Moraceae)	Kafal	Good
<i>Morus alba</i> L. (Moraceae)	Kimbu	Good
<i>Myrica esculenta</i> Buch.- Ham. ex D.Don (Myricaceae)	Ban Kaphal	Fair
<i>Premna bengalensis</i> C.B. Clarke (Verbenaceae)	Kalo Geeneri	Best
<i>Premna latifolia</i> Roxb. Var. Mucronata (Verbenaceae)	Seto Geeneri	Best
<i>Prunus cerasoides</i> D. Don (Rosaceae)	Painyu	Good
<i>Quercus semecarpifolia</i> Smith (Fagaceae)	Kharsu	Best
<i>Rhus javanica</i> L. (Anacardiaceae)	Bhakimlo	Fail
<i>Schefflera venulosa</i> Weight and Arn. Harm. (Araliaceae)	Kutsimal	Good
<i>Toona ciliata</i> M. Roem. (Meliaceae)	Tooni	Fair

Table 5: Wild edible Plants used by the resident of Panchamul, Andhikhola, Syangja

Botanical Name (Family)	Vernacular Name	Parts Used-For
<i>Amaranthus spinosus</i> L. (Amaranthaceae)	Lunde (Herb)	L- Veg
<i>Artocarpus lacucha</i> Buch.-Ham. (Moraceae)	Badahar (Tree)	F-Fruit, Veg
<i>Bambusa bambos</i> (L.) Voss. (Poaceae)	Bans (Bamboo)	Young Shoot
<i>Berberis asiatica</i> Roxb. ex DC. (Berberidaceae)	Chutro (Shrub)	Berry-Fruit
<i>Castanopsis indica</i> (Roxb. Ex Lindl.) A.DC (Fagaceae)	Katoosh (Tree)	Nut-fruit
<i>Centella asiatica</i> (L.) Urb. (Apiaceae)	Ghodtapre (Herb)	L- Pic
<i>Chenopodium album</i> L. (Chenopodiaceae)	Bethe (Herb)	L- Veg
<i>Choerospondias axillaris</i> (Roxb.) (Anacardiaceae)	Lapsi (Tree)	F-Fruit, Jam, Pic
<i>Cinnamomum tamala</i> Buch.- Ham. T.N.& E.(Lauraceae)	Tejpatta (Tree)	L- Spi
<i>Colocasia esculenta</i> (L.) Schott. (Araceae)	Pindalu (Herb)	Root/Tuber- Veg
<i>Crateva unilocularis</i> Buch.-Ham. (Capparaceae)	Siplikaan (Tree)	Shoot,L,F-Veg
<i>Curcuma aromatica</i> Salisb. (Zingiberaceae)	Ban Haledo (Herb)	Rh- Spi
<i>Dendrocalamus hamiltonii</i> Nees & Arn ex M. (Poaceae)	Tama (Bamboo)	Shoot- Veg, Pic
<i>Dioscorea alata</i> L. (Dioscoreaceae)	GharTarul(Climber)	Tuber- Boil, Veg
<i>Dioscorea bulbifera</i> L. (Dioscoreaceae)	Githa (Climber)	Tu/Ro-Boil, Veg
<i>Dioscorea esculenta</i> Lour. (Dioscoreaceae)	Bantarul(Climber)	Root/tuber-Boil, Veg
<i>Dioscorea hamiltonii</i> Hook.f. (Dioscoreaceae)	Bantarul (Climber)	Tu/Ro-Boil, Veg
<i>Dioscorea oppositifolia</i> L. (Dioscoreaceae)	Ban tarul (Climber)	Tuber- Boil
<i>Drymaria cordata</i> (L.) Willd. Ex Schult. (Caryophyllaceae)	Abhijalo (Herb)	L- Veg
<i>Euphorbia royleana</i> Boiss. (Euphorbiaceae)	Saudi (Shrub)	F- Veg
<i>Ficus benghalensis</i> L. (Moraceae)	Bar (Tree)	F-Fruit
<i>Ficus glaberrima</i> Blume (Moraceae)	Pakhri (Tree)	F-Fruit
<i>Ficus sarmentosa</i> Buch.-Ham. Ex Sm. (Moraceae)	Ban timilo (Tree)	Fig-Fruit
<i>Ficus semicordata</i> Buch.-Ham. ex Sm. (Moraceae)	Khaneyu (Tree)	F- Ripe fruit
<i>Fragaria nubicola</i> Lindl. Ex Hook.f. (Rosaceae)	Bhui Aiselu (Shrub)	Berry- Fruit
<i>Justicia adhatoda</i> L. (Acanthaceae)	Asuro (Shrub)	F,L,Fl- Veg, Pic
<i>Lantana camara</i> L. (Verbenaceae)	Ban Phanda (Shrub)	F- Fruit
<i>Lecanthus peduncularis</i> Wall. ex R.) Wedd. (Urticaceae)	Khole (Herb)	L-Veg
<i>Manihot esculenta</i> Crantz (Euphorbiaceae)	Simal Tarul (Shrub)	Tuber- Boil
<i>Mentha spicata</i> L. (Lamiaceae)	Pudina (Herb)	L- Pic
<i>Morus alba</i> L. (Moraceae)	Kafal (Tree)	Berry-Fruit
<i>Myrica esculenta</i> Buch.-Ham. Ex D. Don (Myricaceae)	Ban Kafal (Tree)	F-Fruit, Pic
<i>Nephrolepis cordifolia</i> (L.) C. Prest (Nephrolepidaceae)	Pani amala (Herb)	Tuber-Root
<i>Phyllanthus emblica</i> L. (Phyllanthaceae)	Amla (Tree)	F- Fruit, Pic
<i>Prunus cerasoides</i> D. Don (Rosaceae)	Paiyun (Tree)	F-Ripe Fruit
<i>Psidium guajava</i> L. (Myrtaceae)	Amba (Tree)	F-Fruit
<i>Rhododendron arboreum</i> Sm. (Ericaceae)	Laligurash (Tree)	Juice, Jam
<i>Rubus barberi</i> H.E.Weber (Rosaceae)	Ainselu (Shrub)	Berry-Fruit
<i>Rumex nepalensis</i> Spreng. (Polygonaceae)	Halhale (Herb)	L-Veg
<i>Urtica dioica</i> L. (Urticaceae)	Sisnu (Herb)	L- Veg
<i>Zanthoxylum armatum</i> DC. (Rutaceae)	Timur (Shrub)	F-Fruit, Pic, Spi
L=Leaves, F=Fruit, Veg= Vegetable, Pic=Pickle, Tu/Ro=Tuber/Root, Rh=Rhizome, Spi=Spices, Fl=Flower		

The correlation between plant use value (UV) and the highest fidelity level (%) in the categorized ailments was found significant (Spearman's Correlation test; $r_s = 0.493$, $p(2\text{-tailed}) = 5E-05$) which indicated that by normal standards, the association between use-value (UV) and fidelity level (%) are necessarily those used commonly by the indigenous people of the region.

31 Fodder plant species under 23 genera belonging to 15 families were documented in the study area. Among these Moraceae (8) had the maximum number of species followed by Leguminosae (5), Araliaceae (2), Gramineae (2), Loganiaceae (2), Verbenaceae (2), and the rest with single species each (Fig 4). Out of them, 12 species were categorized as best fodder plants, 09 species as good, and 10 species as fair based on farmer priority and preferences (Table 4). The research reported by (Dhungana et al., 2012; Samant et al., 2007; Upreti and Shrestha, 2006; Panthi, 2003, 2013, Sharma et al., 2016a) were also in close conformity to this study.




Fig 4: Family description of fodder plants in Panchamul




Fig 5: Family description of wild edible plants in Panchamul




Fig 6: Parts of wild edible plants used in Panchamul




Fig 7: Wild edible plants mainly used for in Panchamul

Also, this research has listed 41 wild edible plants belonging to 34 genera and 27 families (Table 5). Of the total families, the dominant is Moraceae (6 Spp.) followed by Dioscoreaceae (5 Spp.), Rosaceae (3 Spp.), Poaceae, Euphorbiaceae, & Urticaceae (2 spp. each), and rest 21 families have one species each (Fig. 5). Wild edible plants used in Panchamul were mainly for fruits (16) followed by leaves (11), tuber (06), root (05), berry (03), rhizome, nuts, and fig (01 each) (Fig. 6). Also, it is estimated that wild edible plants were mainly used for raw fruits (16), followed by vegetables (13), pickles (8) boil (5), spices (3), and jam (2) (Fig. 7). Similar findings were reported in the research by (Bajracharya 1980; Dangol and Gurung 2000; Dangol 2002, 2010; Joshi et al 2007; Acharya and Acharya, 2010; Ghimireya et al 2010; Uprety et al 2012; Shrestha 2013; Dangol et al 2014; Sharma et al., 2016b; Sharma et al., 2017; Khakhalary and Sharma, 2017).

4. CONCLUSION

Panchamul-1 of Andhikhola Rural Municipality is very rich in Non-Timber Forest Products, but more in-depth research exploration is needed. Like, studies on the phytochemical exploration of medicinal plants and their phytotherapeutic sheds of evidence. Though the study area has the best kind of fodder species and numerous wild edible plants, unsustainable harvesting of such plants may cause serious scarcity of fodder and depletion of wild edible plants in the region. It is thus recommended that especially for the most important plant species, a proper cultivation technique should be formulated and conservation and management perspectives should be adopted. Also, medicinal plants with low use-value and fidelity levels should not be overlooked which may lead to the gradual loss of indigenous knowledge. Thus, NTFPs in rural communities should be looked at as a means of livelihood and source of income through sustainable management of the same.

Conflict of interest

The authors declare that they have no conflict of interest.

Funding

There are no funding sources for this paper.

Data and materials availability

All data associated with this study are present in the paper.

REFERENCES AND NOTES

1. Acharya, K.P. and Acharya, R. (2010). Eating from the wild: Indigenous knowledge on wild edible plants in Parroha VDC of Rupandehi district, Central Nepal. *International Journal of Social Forestry (IJSF)* 3(1):28-48.
2. Adhikari, M., Thapa, R., Kunwar, R.M., Devkota, H.p., and Poudel, P. (2019). Ethnomedicinal Uses of Plant Resources in the Machhapuchchhre Rural Municipality of Kaski

- District, Nepal. *Medicines*, 6, 69; doi: 10.3390/medicines6020069.
3. Akanni, K.A. (2013). Economic Benefits of Non-Timber Forest Products among Rural Communities in Nigeria. *Environment and Natural Resources Research*; Vol. 3, No. 4; 2013; <http://dx.doi.org/10.5539/enrr.v3n4p19>.
 4. Arnold, J.E.M., and Perez, M.R. (2001). 'Can non-timber forest products match tropical forest conservation and development objectives?' *Ecological Economics*, 39, 437.
 5. Bajracharya, D. (1980). Nutritive values of Nepalese edible wild fruits. *Z. Lebensm. Unters. Forsch.* 171:363-366.
 6. Baral, S.R. and Kurmi, P.P. (2006) A Compendium of Medicinal Plants in Nepal. Mrs. Rachana Sharma, Kathmandu.
 7. Bhardwaj, A., and Misra, K. (2018). Chapter 11 - Homeopathic Remedies, Editor(s): Kshipra Misra, Priyanka Sharma, Anuja Bhardwaj, Management of High Altitude Pathophysiology, Academic Press, Pages 217-229, ISBN 9780128139998, <https://doi.org/10.1016/B978-0-12-813999-8.00011-2>.
 8. Bhattarai, S.; Chaudhary, R.P.; Taylor, R.S. (2009). Ethno-medicinal Plants Used by the People of Nawalparasi District, Central Nepal. *Our Nat.*, 7, 82–89.
 9. Boullard, B. (2001). Medicinal plants of the world, beliefs and realities, De Boeck Secundair, (ISBN 9782843711176).
 10. Bussmann, R.W., Sharon, D. (2006). Traditional medicinal plant use in Northern Peru: tracking 2000 years of healing culture. *Journal of Ethnobiology and Ethnomedicine* 2, 47.
 11. Camps, F., Claveria, E., Coll, J., Marco, M.P., Messeguer, J., Mela, E. (1990). Ecdysteroid production in tissue cultures of *Polypodium vulgare*, *Phytochemistry*, Volume 29, Issue 12, Pages 3819-3821, [https://doi.org/10.1016/0031-9422\(90\)85339-H](https://doi.org/10.1016/0031-9422(90)85339-H).
 12. Chang, H.C., Agrawal, D.C., Kuo, C.L., Wen, J.L., Chen, C.C., and Tsay, H.S. (2007). "In Vitro Culture of *Drynaria fortunei*, a Fern Species Source of Chinese Medicine "Gu-Sui-Bu"". *In Vitro Cellular & Developmental Biology – Plant*. Springer. 43 (2): 133–139.
 13. Chaudhary, R.P. (1998). Biodiversity in Nepal status and conservation. S. Devi, Saharanpur, India and Craftsman Press, Bangkok, Thailand
 14. Chaudhary, R.P., Upreti, Y. and Joshi, S.P. (2016). Plant biodiversity in Nepal: Conservation and legal status. In *Frontiers of Botany*, Jha, P.K., M. Siwakoti, and S. Rajbhandary(eds.). Central Department of Botany, Tribhuvan University, Kirtipur, Kathmandu, Nepal. pp. 224-268.
 15. Choephel, K. (1993). Dudtsi Mengi Trungpe. Bhojong Mimang Petun Khang, Lhasa. (in Tibetan).
 16. Cochran, W.G. (1977) Sampling Techniques. 3rd Edition, John Wiley & Sons, New York.
 17. Cunningham, A.B. (2001). Applied Ethnobotany: people, wild plant use, and conservation. *Earthscan*, London
 18. Dangol, D., and Gurung, S. B. (2000). Ethnobotanical study of Darai tribe in Chitwan, Nepal. Third National Conference on Science and Technology, Proceedings Vol II, March 8-11, 1999, Royal Nepal Academy of Science and Technology, Kathmandu, pp.1194-1213.
 19. Dangol, D.R. (2002). A final report on on Tharu Ethnobotany Project Part I: Documentation and Conservation of Traditional Knowledge. Submitted to Nepal Forum for Environmental Journalists, Environment Support Fund, Biratnagar, Morang, Nepal.
 20. Dangol, D.R. (2008). Traditional uses of plants of common land habitats in western Chitwan, Nepal. *Journal of the institute of agriculture and animal science*, 29, 71–78.
 21. Dangol, D.R. (2010). Ethnobotany of weeds of Chitwan district, Nepal. *Journal of Natural History Museum* 25:42-53.
 22. Dangol, D.R. Gautam, B. and Oli, B.B. (2014). Wetland plants and their local uses: Observations from Rampur ghol, Chitwan, Nepal. *Journal of Natural History Museum* 28:142-159
 23. Dawa, D. (1999). A Clear Mirror of Tibetan Medicinal Plants. Tibet Domani, Rome, Italy.
 24. Dhungana, S., Tripathee, H.P., Puri, L., Timilsina, Y.P., and Devkota, K.P. (2012). Nutritional analysis of locally preferred fodder trees of middle hills of Nepal: A Case Study from Hemja VDC, Kaski District. *Nepal J. Sc. and Tech.* 13(2):39-44.
 25. DPR. (2001). Flowering Plants of Nepal (Phanerogams), His Majesty's Government, Ministry of Forests and Soil Conservation, Department of Plant Resources, Kathmandu, Nepal.
 26. Duvey, B.K., Goyel, R. Parashar, B., Verma, D., Dhameja, H., Sharma, D. (2012). *Trichosanthes tricuspidata*: Exploration of Its Medicinal Value. *Asian J. Pharm. Tech.* 2(1): Page 26-28.
 27. Edwards, D.M., and M.R. Maharjan. (1994). Non-Timber Forest Products in Nepal: the potential for community management. Nepal UK Community Forestry Project Report. 6pp.
 28. Food and Agriculture Organization (FAO) (1999). FAO Forestry – Towards a Harmonised Definition of Non-Wood Forest Products. *Unasylva*– No. 198, Vol. 50
 29. Food and Agriculture Organization of the United Nations (FAO) (1995). Non-Wood Forest Products for Rural Income and Sustainable Forestry. Rome: FAO NWFPs 7.
 30. Friedman, J., Yaniv, Z., Dafni, A., Palewitch, D. (1986). A preliminary classification of the healing potential of medicinal plants, based on a rational analysis of an ethnopharmacological field survey among Bedouins in the Negev desert, Israe. *Journal of Ethnopharmacology* 16, 275–287.

31. Gazzaneo, L.R.S., Lucena, R.F.P., Albuquerque, U.P. (2005). Knowledge and use of medicinal plants by local specialists in an region of Atlantic forest in the state of Pernambuco (Northeastern Brazil). *Journal of Ethnobiology and Ethnomedicine* 1, doi:10.1186/1746-4269-r1-9
32. Ghimeray, A., Sharma, P., Ghimire, B., Lamsal, K., Ghimire, B.K., and Cho, D.H. (2010). Wild edible angiospermic plants of Ilam Hills (Eastern Nepal) and their mode of use by the local community. *Korean J. Pl. Tax.* 40(1):1-4.
33. Ghimire, S.K., D.B. Parajuli, T.N. Gurung, and Y.C. Lama (1999). Conservation of Plant Resources, Community Development, and Training in Applied Ethnobotany at Shey-Phoksundo National Park and its Buffer-zone, Dolpa. WWF Nepal Program Report Series No. 38, WWF Nepal Program, Kathmandu, Nepal.
34. Ghimire, S.K., Lama, Y.C., Tripathi, G.R., Schmitt, S., Aumeeruddy-Thomas, Y. (2001). Conservation of plant resources, community development, and training in applied ethnobotany at Shey-Phoksundo national park and its buffer zone, Dolpa. Report Series No. 41, WWF Nepal Program, Kathmandu, Nepal.
35. Ghimire, S.K., Y.C. Lama, Amchi T.N. Gurung, and Y. Aumeeruddy Thomas (2000). Conservation of Plant Resources, Community Development, and Training in Applied Ethnobotany at Shey-Phoksundo National Park and its Buffer zone, Dolpa. Third Year. WWF Nepal Program Report Series No. 40, WWF Nepal Program, Kathmandu, Nepal.
36. Golam, R., Madhav, K., Ram, P.S. (2008). The role of non-timber forest products in poverty reduction in India: Prospects and problems. *Development in Practice*, 6: 779 – 788.
37. Goyal, R. K., Singh, J., Lal, H. (2003). "*Asparagus racemosus*—an update". *Indian Journal of Medical Sciences*. 57 (9): 408–414. PMID 14515032.
38. Gurav, S.S., Gurav, S.N. (2014). A Comprehensive Review: *Bergenia ligulata* Wall - A Controversial Clinical Candidate. *International Journal of Pharmaceutical Sciences and Research*. DOI: [http://dx.doi.org/10.13040/IJPSR.0975-8232.5\(5\).1630-42](http://dx.doi.org/10.13040/IJPSR.0975-8232.5(5).1630-42).
39. Hara, H., and Williams, L.H.J. (1979). Enumeration of flowering plants of Nepal. Vol. 2. British Museum (Natural History) London.
40. Hara, H., Chater, A.O. and Williams, L.H. J. (1982). An enumeration of flowering plants of Nepal. Vol. 3. British Museum (Natural History) London.
41. Hara, H., Stearn, W.T. and Williams, L.H.J. (1978). An enumeration of flowering plants of Nepal. Vol. 1. British Museum (Natural History) London.
42. Hasan, A. (2018). Medicinal importance of prickly pear cactus. 4th World Congress on Medicinal Plants & Natural Products Research and 12th Global Ethnomedicine & Ethnopharmacology Conference, Osaka, Japan, *Medicinal & Aromatic Plants*, Volume 7 DOI: 10.4172/2167-0412-C1-017.
43. Hasan, M.K., Gatto, P., and Jha, P.K. (2013). Traditional uses of wild medicinal plants and their management practices in Nepal-A study in Makawanpur district. *Int. J. Med. Arom. Plants*, ISSN 2249 – 4340. Vol. 3, No. 1, pp. 102-112.
44. Hechtman, L. (2018). *Clinical Naturopathic Medicine* (2ed.). Elsevier. pp. 879, 908. ISBN 9780729542425.
45. Heinrich, M., Ankli, A., Frei, B., Weimann, C., Sticher, O. (1998). Medicinal plants in Mexico: healers' consensus and cultural importance. *Social Science and Medicine* 47, 91–112.
46. Jain, S.K., Saklani, A. (1991). Observations on the ethnobotany of the Tons valley region in the Uttarkashi district of the Northwest Himalaya, India. *Mountain Research and Development* 11, 157–161.
47. Joshi, A.R., Edington, J.M. (1990). The use of medicinal plants by two village communities in the Central development region of Nepal. *Economic Botany* 44, 71–83.
48. Joshi, K.; Joshi, R.; Joshi, A.R. (2011). Indigenous knowledge and uses of medicinal plants in Macchegaun Nepal. *IJTK*, 10, 281–286.
49. Joshi, K.K., Joshi, S.D. (2001). Genetic Heritage of Medicinal and Aromatic Plants of Nepal Himalaya. Buddha Academic Publishers and Distributors Pvt. Ltd., Kathmandu, Nepal
50. Joshi, N., Kehlenbeck, K. and Maass, B.L. (2007). Traditional, neglected vegetables of Nepal: Their sustainable utilization for meeting human needs. Conference on International Agricultural Research for Development, Tropentag, University of Kassel-Witzenhausen and University of Gottingen, 9-11 Oct 2007.
51. Kapil A, Sharma S. (1997). Immunopotentiating compounds from *Tinospora cordifolia*. *J Ethnopharmacol.* 1997; 58(2):89-95.9406896.
52. Karki, S. (1996). Investigating non -timber forest products (NTFPs) opportunities in Nepal. Nepal Australia Community Forestry Project. 16p.
53. Khakhalary, B., and Sharma, S. (2017). Non-Timber Forest Product: Case Study Of Diversity In Garampani Wildlife Sanctuary, KarbiAnglong District, Assam, India. *International Journal of Innovative Research and Advanced Studies (IJIRAS)*. Vol.4 (5). Pp 53-55.
54. Kremen, C., and Merenlender, M. (2018). Landscapes that work for biodiversity and people, *Science*, 362, pp. 1-9, 10.1126/science.aau6020
55. Kumar A., Rana A.C. (2013) Pharmacognostic and pharmacological profile of traditional medicinal plant: *Myrica nagi*. *Int. Res. J. Pharm.* 2013; 3:32–37.

56. Kumar, D., Kumar R., and Sharda, K. (2018). "Medicinal property of Nirgundi" *Journal of Pharmacognosy and Phytochemistry*. SP1: 2147-2151.
57. Kunwar, R.M., Nepal, B.K., Kshhetri, H.B., Rai, S.K., Bussmann, R.W. (2006). Ethnomedicine in Himalaya: a case study from Dolpa, Humla, Jumla and Mustang districts of Nepal. *Journal of Ethnobiology and Ethnomedicine* 2, 27.
58. Lama, Y.C., Ghimire, S.K., Aumeeruddy-Thomas, A. (2001). Medicinal Plants of Dolpo: Amchis' Knowledge and Conservation. People and Plants Initiative and WWF Nepal Program, Nepal.
59. Luitel, D.R., Rokaya, M.B., Timsina, B., Munzbergová, Z. (2014). Medicinal plants used by the Tamang community in the Makawanpur district of Central Nepal. *J. Ethnobiol. Ethnomed.* 10, 5.
60. Malla, B.; Chhetri, R.B. (2009). Indigenous knowledge on ethnobotanical plants of Kavrepalanchowk District. *Kathmandu Univ. J. Sci. Eng. Technol.* 5, 96–109.
61. Manandhar, N.P. (2002). Plants and People of Nepal. Timber Press Inc., Portland, Oregon.
62. Martin, G.J. (1995). Ethnobotany. A People and Plants conservation manual. Chapman and Hall 447.
63. MoEF (2018). Nepal's Sixth National Report to the Convention on Biological Diversity. Ministry of Forests and Environment, Government of Nepal.
64. NEHHPA (Nepal Herbs and Herbal Products Association). https://www.nehha.org.np/herbs_directory/armelia-nepalensis-tayl.
65. Nguyen, D.D., Nguyen, N.H., Nguyen, T.T., Phan, T.S., Van Nguyen, D., Grabe, M., Johansson, R., Lindgren, G., Stjernstrom, N.E., and Oderberg, T.A. (1996) The use of water extracts from the bark of *Choerospondias axillaris* in the treatment of second degree burns. *Journal of Plastic Surgery and Hand Surgery* 30: 139.
66. NPHC (2011). National Population and Housing Census 2011, Central Bureau of Statistics, Government of Nepal, Kathmandu, Nepal. Kathmandu, Nepal: CBS.
67. Panging, S.M., Sharma, S. (2017). Studies on Ethno Medicinal and Traditional Healing Practices among Mising Community of Desangmukh Gaon Panchayat, Sivasagar District of Assam, India. *Journal of Medicinal Plants Studies*; 5(4): 193-196. Baral, S.R., Kurmi, P.P. (2006). A Compendium of Medicinal Plants in Nepal. Mrs. Rachana Sharma, Kathmandu, Nepal.
68. Panthi, M.P. (2003). Fodder plants and their indigenous management in Adguri and Narapani Village Development Committees (VDCs), West Nepal. A research report submitted to the Dean's office, Institute of Science and Technology, Tribhuvan University, Kathmandu.
69. Panthi, M.P. (2003). Indigenous knowledge on use of local fodder trees in mid hills of west Nepal. *Tribhuvan University Journal*, Volume. Xxviii, Numbers. 1-2, pp 171-180.
70. Patil, D., Roy, S., Dahake, R., Rajopadhye, S., Kothari, S., Deshmukh, R., Chowdhary, A. (2013). Evaluation of *Jatropha curcas* Linn. leaf extracts for its cytotoxicity and potential to inhibit hemagglutinin protein of influenza virus. *Indian J Virol*; 24(2):220-6. doi: 10.1007/s13337-013-0154-z.
71. Peters, C.M., Gentry, A.H., and Mendelsohn, R.O. (1989). Valuation of the Amazonian rainforest, *Nature*, 339(29), June, 655.656.
72. Phillips, O., Gentry, A.H., Reynel, C., Wilkin, P., Galvez Durand, B.C. (1994). Quantitative ethnobotany and Amazonian conservation. *Conservation Biology* 8, 225–248.
73. Phole, P. (1990). Useful Plants of Manang District: A Contribution to the Ethnobotany of the Nepal Himalaya. Franz Steiner Verlag, Stuttgart.
74. Pizzorno Jr., Joseph E.; Murray, M.T.; Joiner-Bey, H. (2015). Clinician's Handbook of Natural Medicine (3rd ed.). Churchill Livingstone. p. 516. ISBN 9780702055140.
75. Polunin, O., and Stainton, A. (1984). Flowers of the Himalaya. Oxford University Press. India.
76. Press, J.R., Shrestha, K.K. and Sutton, D.A. (2000). Annotated checklist of the flowering plants of Nepal. The Natural history Museum, London.
77. Rajbhandari, K.R. (2001). Ethnobotany of Nepal. Ethnobotanical Society of Nepal, Kathmandu, Nepal.
78. Rajkumar, N., Shivanna, M.B. (2009). Ethno-medicinal application of plants in the Eastern region of Shimoga district, Karnataka, India. *Journal of Ethnopharmacology* 126, 64–73.
79. Rajkumar, N., Shivanna, M.B. (2009). Ethno-medicinal application of plants in the Eastern region of Shimoga district, Karnataka, India. *Journal of Ethnopharmacology* 126, 64–73.
80. Rastogi, A., Godbole, A., and Shengji, P. (1998). Applied ethnobotany in natural resource management- traditional home gardens. International Centre for Integrated Mountain Development (ICIMOD), Kathmandu, Nepal.
81. Rawat, P., Rai, N., Kumar, N., Bachheti, R.K. (2017). Review on *Rhododendron arboretum* - a magical tree. *Orient Pharm Exp Med* 17, 297–308. <https://doi.org/10.1007/s13596-017-0289-3>.
82. Rokaya, M.B., Münzbergová, Z., and Timsina, B. (2010). Ethnobotanical study of medicinal plants from the Humla district of western Nepal. *Journal of Ethnopharmacology* 130, 485–504. doi:10.1016/j.jep.2010.05.036.
83. Ros-Tonen, M.A.F. (2000). The role of non-timber forest products in sustainable tropical forest management. *Holz als Roh- und Werkstoff* 58: 196-201.

84. Rysdyk, E. C. (2019). *The Nepalese Shamanic Path: Practices for Negotiating the Spirit World*. Simon and Schuster. ISBN 978-1-62055-795-2.
85. Samant, S.S., Singh, M., Lal, M. and Pant, S. (2007). Diversity, distribution and prioritization of fodder species for conservation in Kullu District, Northwestern Himalaya, India. *J. of Mountain Science* 4(3): 259-274.
86. Sayer, J. (2009). Reconciling Conservation and Development: Are Landscapes the Answer? *Biotropica*, 41: 649-652. doi:10.1111/j.1744-7429.2009.00575.x
87. SCBD (2001). 'The Value of Forest Ecosystem', CBD Technical Series No.4, Secretariat of the Convention on Biological Diversity, Montreal, Canada.
88. Sharma, P. (1996). Non-wood forest products and integrated mountain development: Observations from Nepal. Business Seminar on Medicinal Herbs, Essential Oils, and Other Non-Timber Forest Products, held in Kathmandu, December 1996. DEG/NGCCI. 11pp.
89. Sharma, S. Choudhury, P., Chetry, N. (2016a). Effect of Hill Cutting on the Diversity of Fodder Plants in the Raid-Marwet Region, Ri-Bhoi District of Meghalaya. In *Conserving Biological Diversity: A Multiscaled Approach* (Chapter: 5) Research India Publication (Ed.) Aabid Hussain Mir and Nazir Ahmad Bhat. pp 65-78.
90. Sharma, S., Choudhury, P and Chetry, N. (2016b). Diversity of Non-Timber Forest Products (NTFPs): A Provisioning Ecosystem Services among the Marwet Community, Ri-Bhoi District, Meghalaya. *International Journal of Scientific and Research Publication*, 6(9), 48-55.
91. Sharma, S., Choudhury, P and Chetry, N. (2017). Land Degradation and its Impacts on some major Ecosystem Services in Raid-Marwet Region, Ri-Bhoi District of Meghalaya, India. Ph.d. Thesis, Department of Ecology and Environmental Science, Assam University, Silchar. <http://hdl.handle.net/10603/291889>
92. Shiva, M.P., Verma, S.K. (2002). *Approaches to Sustainable Forest Management and Biodiversity Conservation: With Pivotal Role of Non-timber Forest Products*. Dehra Dun: Centre for Minor Forest Products, Valley Offset Printers.
93. Shrestha, D. (2013). Indigenous vegetables of Nepal for biodiversity and food security. *International Journal of Biodiversity and Conservation* 5(3):98-108.
94. Shrestha, P.M. & Dhillon, S.S. (2003). Medicinal plant diversity and use in the highlands of Dolakha district, Nepal. *J. Ethnopharmacol*, 86, 81-96.
95. Shrestha, T.B., and Joshi, R. M. (1996). *Rare, endemic, and endangered plants of Nepal*. WWF Nepal Program, Kathmandu, Nepal
96. Sood P., Shri R. (2018). A review on ethnomedicinal, phytochemical, and pharmacological aspects of *Myrica esculenta*. *Indian J. Pharm. Sci.*; 80:2-13.
97. Srithi, K., Balslevb, H., Wangpakapattanawonga, P., Srisangac, P., Trisonthia, C. (2009). Medicinal plant knowledge and its erosion among the Mien (Yao) in northern Thailand. *Journal of Ethnopharmacology* 123, 335-342.
98. Stainton, A. (1988). *Flowers of Himalaya: a supplement* oxford University Press, India.
99. Subedi, B.P., Bhattarai, N.K. (1998). Community managed enterprise: participation of rural people in medicinal and aromatic plants conservation and use. In: *A Paper Presented at International Conference on Medicinal Plants*, February 16-20, 1998, Bangalore, India.
100. Subedi, B.P. (1997). Utilization of non-timber forest products: issues and strategies to environmental conservation and economic development. Theme paper presented in the workshop on —The utilization of NTFPs for environmental Conservation and Economic Development in Nepal organized by ANSAB on March 29, 1997, in Kathmandu.
101. Subedi, B.P., Ghimire, P.L., Koontz, A., Khanal, S.C., Katwal, P., Sthapit, K.R., and Mishra, S.K. (2014). Private Sector Involvement and Investment in Nepal's Forestry: Status, Prospects, and Way Forward. Multi-Stakeholder Forestry Programme (MSFP)- Service Support Unit, Babarmahal, Kathmandu, Nepal.
102. Tamang, G. (2003). An ethnobiological study of Tamang people. *Our Nat.* 1, 37-41.
103. Tamang, R.; Sedai, D.R. (2016). Documentation of Ethnomedicinal Knowledge on Plant Resources Used by Baram Community in Arupokhari VDC, Gorkha District, Central Nepal. *Bull. Dept. Plant Res.*, 38, 60-64.
104. Thapa, S. (2012). Medico-ethnobotany of Magar Community in Salija VDC Parbat district, Central Nepal. *Our Nat.* 10, 176-190.
105. Thapa, S. (2012). Medico-ethnobotany of Magar Community in Salija VDC Parbat district, Central Nepal. *Our Nat.* 2012, 10, 176-190.
106. Trotter, R.T., and Logan, M.H. (1986). Informant consensus: a new approach for identifying potentially effective medicinal plants. In: Etkin, N.L. (Ed.), *Plants in Indigenous Medicine and Diet*. Bedford Hills, New York. Redgrave Publishing Company, pp. 91-112
107. Umberto, Q. (2012). *CRC World dictionary of medicinal and poisonous plants: common names, scientific names, eponyms, synonyms, and etymology*. Volume 5 R-Z. CRC Press Taylor and Francis Group, page 228.
108. Upreti, C.R. and Shrestha, B.K. (2006). Nutrient contents of feeds and fodder in Nepal. Animal Nutrition Division, NARC, Khumaltar, Lalitpur.
109. Upreti, Y., Asselin, H., Boon, E.K., Yadav, S., Shrestha, K.K. (2010). Indigenous use and bio-efficacy of medicinal plants

- in the Rasuwa district, Central Nepal. *Journal of Ethnobiology and Ethnomedicine* 6, 3.
110. Uprety, Y., Poudel, R.C., Shrestha, K.K., Rajbhandary, S., Tiwari, N.N., Shrestha, U.B. and Asselin, H. (2012). Diversity of use and local knowledge of wild edible plant resources in Nepal. *Journal of Ethnobiology and Ethnomedicine* 8:16. DOI: <http://dx.doi.org/10.1186/1746-4269-8-16>.
111. Wan Hassan, W.E. (2010). *Ulam: Salad Herbs of Malaysia*. Masbe Sdn. Bhd. pp. 106–107. ISBN 9789834466404.
112. Wang, H., Gao, X.D., Zhou, G.C., Cai, L., and Yao, W.B. (2008) In vitro and in vivo antioxidant activity of aqueous extract from *Choerospondias axillaris* fruit. *Food Chemistry* 106: 888–895.
113. Wong, J. (2002). Biometrics and NTFP Inventory. School of Agricultural and Forest Sciences, University of Wales, Bangor, Gwynedd LL57 2UW, UK.
114. Yang, L., and Jiang, J-G. (2009). Bioactive components and functional properties of *Hottuynia cordata* and its applications, *Pharmaceutical Biology*, 47:12, 1154-1161, DOI: 10.3109/13880200903019200.
115. Zheng-yi, W and P.H. Raveen (ed.), 1994. Flora of China. Science press (Beijing) and Missouri Botanical Garden, USA.